



The HerBS sample

Finding lenses in the wide-area, sub-mm
H-ATLAS survey with SCUBA-2 observations





Gravitational lenses provide otherwise unreachable levels of detail



ALMA Partnership 2015, Dye et al. 2015

Gravitational lenses provide otherwise unreachable levels of detail




Dye et al. 2015

We use the 590 sq. deg. H-ATLAS survey to look for lenses

H-ATLAS:



590 sq. deg.
NGP, SGP & GAMA
Confusion limits

**Data Release
TODAY**
www.h-atlas.org




Eales et al. 2009, Valiante et al. 2016

Gravitational lenses provide otherwise unreachable levels of detail



Dye et al. 2015

Large surveys can't directly distinguish lenses





The HerBS sample contains lensed ULIRGs and unlensed HyLIRGs

HerBS selection:

236 sources with

$z_{\text{phot}} > 2$




$S_{500\mu\text{m}} > 80 \text{ mJy}$



Negrello et al. 2009

Pearson et al. 2013


Source confusion effects need to be measured

	<i>Herschel</i>		
λ [μm]	250	350	500
Angular size	18"	25"	36"
Surface	158%	306%	634%
Beam size			


A lensed sample that probes the Cosmic Star Formation History

Of our 189 SCUBA-2 observed galaxies, we expect 150 galaxies to be lensed


Sample purification



HerBS sample



Lensing estimates



Our sample contained several local galaxies and QSO's

Herschel/SPIRE
250 μm

Herschel/SPIRE
350 μm

Herschel/SPIRE
500 μm


JCMT/SCUBA-2
850 μm

SED
SPIRE - SCUBA-2


**Of which only ~5
show extended
profiles**

Template fitted shows little changes to previous version


	Best-fit	Pearson
T_{cold}	19.0 ± 3.3	23.9
T_{hot}	39.7 ± 7.1	46.9
α	46.1 ± 23.1	30.1
β	2.62 ± 0.82	2 (fixed)
χ^2	929	1143




We find no HerBS selection problems




Our sources probe high redshifts and luminosities



Cosmological source count models suggest an 80% lensing fraction




Cai et al. 2013




Preliminary analysis of VIKING data shows at least 70% lenses

No counterpart




8%

Barely resolved



19%

Clear counterpart





73%

Preliminary!!!



A lensed sample that probes the Cosmic Star Formation History

Of our 189 SCUBA-2 observed galaxies, we expect 150 galaxies to be lensed



Lensing estimates



HerBS sample



Hi-res Lenses



The HerBS sample

Finding lenses in the wide-area, sub-mm
H-ATLAS survey with SCUBA-2 observations

